

Effect of Plant Growth Promoting Activities of PG-63 in Salinity Stress on *Trigonella foenum-graecum*

Drashti Patel^{1,2}, Dimple Pardhi¹, Tejalba Rathod¹, Bansi Purohit¹, Daxa Damor², Rushikesh Joshi², Kiransinh Rajput^{*1}

¹Department of Microbiology and Biotechnology, School of Science, Gujarat University
Ahmedabad-380009, Gujarat, INDIA

²Department of Biochemistry and Forensic Sciences, School of Science, Gujarat University
Ahmedabad-380009, Gujarat, INDIA

Email: rajputkn@gujaratuniversity.ac.in

ABSTRACT

Salt stress is one of the important limiting factor for agricultural productivity of soil and become a significant barrier to crop production globally. Natural preparations like panchagavya (made from cow, dung, urine, milk, ghee) can be used in agricultural practices which increases agricultural productivity and soil fertility. 64 bacteria were isolated from fermented panchagavya and checked their salt tolerance ability in the presence of different salt concentrations. The PG-63 isolate shows better tolerance in salinity stress. In the present study, PG-63 shows tolerance towards salt stress. The PG-63 was further checked for plant growth promoting activities. PG-63 showed 22.67 µg/ml IAA production in control and 25.15 µg/ml in the presence of salt. The highest GA production by PG-63 was 3.38 mg/ml in control and 2.19 mg/ml in the presence of salt. Ammonia production was 36 µmol/ml in the control and 33.8 µmol/ml in the presence of salt. The phosphate solubilization was 64.40 µg/ml in the control and 70.67 µg/ml in the presence of salt. The seed germination index of *Trigonella foenum-graecum* was observed 81 in the presence of salt and PG-63. The treated plants were longer than the untreated control plants in terms of their roots and shoots. On the 15th day, PG-63 treated plants showed an increase in root (12%) and shoot (12.5%) length as well as an increase in leaf length (9.9%).

Keywords: Panchagavya, salinity, indole-3-acetic acid, gibberellic acid, phosphate solubilization

Received 24.05.2023

Revised 11.06.2023

Accepted 11.08.2023

How to cite this article:

Drashti Patel, Dimple Pardhi, Tejalba Rathod, Bansi Purohit, Daxa Damor, Rushikesh Joshi, Kiransinh Rajput. Effect of Plant Growth Promoting Activities of PG-63 in Salinity Stress on *Trigonella foenum-graecum*. Adv. Biores., Special Issue 1: 2023: 160-169

INTRODUCTION

The numerous both abiotic and biotic variables have a significant impact on agricultural practices all over the world. Farming practices are significantly hampered by a number of these abiotic variables. The growth of plants is influenced by abiotic elements such as temperature, soil salinity, drought stress, and heavy metal stress. The salinity of the soil has emerged as one of the major problems limiting crop productivity globally. Due to anthropogenic activity, the rate of soil salinization has accelerated in recent years (23). Crop productivity is lost, and salt levels are high in areas of soil where there is an excessive input of fertilizer (32). The application of fertilizers with a high salt index has an osmotic influence that makes it challenging to access the water required for plant growth (14). It is also claimed that agricultural techniques have an impact on crop productivity on saline soils. In these circumstances, more salt is accumulated because deeper tilling or ploughing of the soil causes more water to evaporate from its surface. Additionally, according to (8), salts in irrigation water may increase the salinity of the soil, making it less able to support plant growth. Most plants in nature are sensitive to salt stress because it may hinder microbial growth in their surrounding habitat and disrupt the normal plant-microbe interaction in soil (1). Salinization has an impact on all phases of plant development, including germination, physiological development, reproductive development, and senescence. Salinity in the soil has a negative effect on the general health of plants. Salinity has an impact on the physicochemical properties of the soil, the ecological balance of the area, and the agricultural production of most crops. Excessive salt consumption reduces leaf size, chlorophyll content, and the efficiency of photosynthetic activities. An essential part of the electron transport chain

(ETC), photosystem II (PSII), is suppressed when subjected to salt stress (25,17). The growth and development of ST-PGPB (salt-tolerant plant growth promoting bacteria) may be crucial in rescuing crops from such stressful situations (20). The natural defence against salinity stress conditions is the presence of microorganisms that promote plant growth (9). Under stressful circumstances, plant-microbe interaction is essential for maintaining soil characteristics, crop productivity, and microbial diversity. The interaction between plants and microorganisms is also crucial for plant growth and development in a dynamic environment. Plant growth-promoting microorganisms (PGPM) are a variety of bacteria that cluster around plant roots and promote the growth and development of the plant when it is exposed to salinity stress (19). According to (30), some halophilic bacteria have been shown to be effective at lowering plant salt stress, which can help to improve the growth and productivity of crops grown in salinity. Numerous studies have shown that bacteria like *Achromobacter*, *Arthrobacter*, *Bacillus*, *Chryseobacterium*, *Enterobacter*, *Ochrobactrum*, and *Pseudomonas*, which have been isolated from saline soil, can facilitate plant growth in saline environments (33, 16, 35, 4). Salinity control and the promotion of plant growth and development can both benefit from PGPB-mediated amelioration of salinity stress. The global trend towards organic farming is expanding. Currently, there are 0.7 million organic farms spread across 30.4 million hectares in more than 130 countries. Panchagavya is an organic formulation, which in Sanskrit means the blend of five cow products i.e., milk, ghee, curd, dung, and urine (all these products are individually called as "Gavya" and collectively named as panchagavya). "Panchagavya" has got reference in the scriptures of Vedas (divine scriptures of Indian wisdom) and Vrikshayurveda (31). They are a plentiful supply of microbial communities, macronutrients, micronutrients, and chemicals that support plant growth, such as immune boosters (21). PG-63 bacterial isolate was obtained from fermented panchagavya. The present study is focused on the characterization of PG-63 bacterial isolate and to check its effect on *Trigonella foenum-graecum* for salt tolerance. PG-63 shows good salt tolerance even at high concentrations of NaCl. In this study, we are aiming to find out the potential roles of PG-63 in salinity stress.

MATERIALS AND METHODS

Isolation of bacterial isolates for fermented panchagavya

64 bacterial cultures were isolated from fermented panchagavya (12). Various plant growth promoting traits like IAA production, GA production, ammonia production, phosphate solubilization, EPS production were checked in salinity stress condition. PG-63 bacterial isolate showed better survival in the presence of NaCl. Results were shown in table-2.

Inoculum preparation

The bacterial cultures from slants were inoculated in sterile nutrient broth and incubated at 30°C for 24 h. Cells were separated by centrifugation at 5000 rpm for 10 min and the supernatant was removed. The cell pellet was resuspended in sterile normal saline to get optical density of 1.0. The prepared culture suspension was used as a 1.0% (v/v) inoculum to study the plant growth-promoting parameters like IAA, GA, NH₃ production and phosphate solubilization.

The absorbance was measured in spectrophotometer at 600nm after 24 hours. The standard calibration curve of BaCl₂ (1.0 O.D ~ 8×10⁸ cells/ml) was prepared to calculate the number of cells.

Salinity stress response of PG-63

The salt tolerance of PG-63 was checked by observing as their growth in the nutrient broth medium supplemented with different NaCl concentrations (1, 3, 5, 7, 10 and 15% w/v). PG-63 was inoculated in sterile nutrient broth and incubated at 28°C for 24 h.

Plant growth promoting traits

The selected bacterial cultures were tested qualitatively and quantitatively for multiple plant growth promoting activities like indole-3-acetic acid production (IAA), gibberellic acid production (GA), ammonia production (NH₃), phosphate solubilization (P).

• Indole-3-acetic acid production

For IAA production, 50.0 ml of Luria-Bertani broth (pH 7.5) containing 0.1% (w/v) L-tryptophan in 250 ml flask was inoculated with 1.0% (v/v) inoculum and incubated at 30°C in the dark (as IAA is light sensitive), 120 rpm on an orbital shaker. IAA production was estimated from samples withdrawn at 24-h intervals up to 24–144 h until production was decreased.

The spectrophotometric estimation of IAA was done according to (5). Culture supernatant (1 ml) was mixed with 2.0 ml of Salkowski reagent and incubated at room temperature (RT) in the dark for 30 min. Development of pink colour shows the production of IAA and absorbance was recorded at 530 nm. The standard calibration curve of IAA (100 µg/ml) was prepared to calculate the IAA production.

• Gibberellic acid production

Gibberellic acid production was carried out in 50.0 ml nutrient broth (pH 7.4) in 250 ml flask, 1.0% (v/v) inoculum, at 30°C, 120 rpm on an orbital shaker for 96 h. GA production was estimated from samples withdrawn at 24 h intervals up to 24–96 h until it was decreased.

An equal volume of cell-free supernatant and ethyl acetate (EA) was taken in a test tube and shook vigorously. The EA extract was collected separately in a glass beaker, and the extraction was repeated three times. The separated EA extract was evaporated by rotary evaporator at 45°C. The residues were dissolved in 2.0 ml methanol. To this, 1.0 ml DNPH (2, 4 – dinitrophenylhydrazine) was added and incubated in a boiling water bath for 5 min. After incubation, it was cooled in an ice-water bath, and 5.0 ml of 10% (w/v) alcoholic potassium hydroxide was added, allowed to stand for 10 min at RT. To this, 15.0 ml of sterile distilled water was added, and the intensity of the color (red wine) produced was measured at 430 nm (11). A standard calibration curve of GA (0.8 mg/ml) was prepared.

- **Ammonia production**

Ammonia production was determined in a 50-ml flask containing 20.0 ml sterile peptone water, pH 7.2. It was inoculated with 1.0% (v/v) inoculum and incubated at 30°C, 120 rpm on an orbital shaker for 144 h. The amount of NH₃ produced was estimated from samples withdrawn at 24 h intervals up to 24–144 h until it was decreased.

Spectrophotometric estimation of ammonia was done as described by (6). Culture supernatant (1 ml) was mixed with 0.1 ml Nessler's reagent, and the final volume was made to 5.0 ml by adding ammonia-free distilled water. NH₃ production is indicated by a change in color from yellow to brown, and absorbance was measured at 425 nm. The NH₃ produced was calculated using the ammonium hydroxide (0.2 μmol/ml) standard calibration curve.

- **Phosphate solubilization**

The qualitative phosphate solubilization test was carried out using spot inoculation technique on an agar plate containing an insoluble tricalcium phosphate (TCP) and incubated at 30°C for 120 h (29). A zone develops surrounding the colony because of phosphate solubilization. The phosphate solubilization index (PSI) was calculated using the formula below.

$$PSI = \frac{\text{Colony diameter} + \text{Zone of clearance}}{\text{Colony diameter}}$$

For quantitative phosphate solubilization, 100 ml nutrient broth in 250 ml flasks were inoculated and incubated at 30°C, at 120 rpm on an orbital shaker for 15 days. The amount of phosphate solubilization was measured at 5 days interval for up to 15 days.

A 2.0 ml sample was centrifuged at 10,000 rpm for 20 min to separate the cells. To 0.1 ml supernatant, 0.9 ml double distilled water and 1.0 ml chloromolybdic acid was added. Contents were diluted by adding 4.0 ml double distilled water. To this, 25.0 μL chlorostannous acid reagent was added, and the test tubes were mixed well till blue color developed. Absorbance was measured at 600 nm. The amount of phosphate solubilized was calculated against K₂HPO₄ (100.0 μg/ml) standard calibration curve.

Seed germination test

For the germination test, seeds were first surface sterilized by gently shaking in 70% methanol for 2 minutes, then the seeds were washed with sterile distilled water. The seeds were dipped in 0.1% (v/v) HgCl₂ for 1 minute and again seeds were washed twice with sterile distilled water. After surface sterilization, the seeds were placed on sterile cotton in a sterile Petri plate. The experiment was performed in 2 sets.

1) *Trigonella foenum-graecum* seeds plus distilled water

2) *Trigonella foenum-graecum* seeds plus NaCl(2%w/v) + PG-63

After 48 hours, germinated seeds were counted and the germination percent, seed vigour index was calculated by the following formula:

$$\text{Seed vigour index} = \text{seedling length} \times \text{germination percent}$$
$$\text{Germination percentage} = \frac{\text{Number of germinated seeds} \times 100}{\text{Number of seeds}}$$

Pot study

Pot study was carried out in plastic bags having 18×18 cm size in the month of February to March 2023, when the average temperature was below 30°C. For preparation of pots, collected farm soil was sun dried for 5 days. Approximately 4.0 kilograms of sun-dried soil was filled in the pots. Salt tolerance of the PG-63 was tested at 1% NaCl. The seeds were surface sterilized by gently shaking them in 70% methanol for 2 minutes, then the seeds were washed with sterile distilled water, next the seeds were dipped in 0.1% HgCl₂; for 1 minute, then the seeds were again washed twice with sterile distilled water. The 10 numbers of seeds

were sown in the pot at 1 cm depth. Irrigation was done with tap water. For preparation of bacterial suspension for pot treatment, the PG-63 bacterial isolate was scraped from preserved slant and inoculated in 100 ml sterile nutrient broth, incubated at 30°C, at 120 rpm for 24 hours. After 20 days, plants were uprooted and vegetative parameters like root length, shoot length, total plant length, wet weight, dry weight was measured. The pot experiment design is shown in table-1.

RESULTS AND DISCUSSION

PG-63 was studied for its salt tolerance. PG-63 shows better growth in the presence of different NaCl (%w/v) concentrations. PG-63 shows plant growth promoting characteristics like indole-3-acetic acid production, gibberellic acid production, ammonia production, phosphate solubilization etc. in the presence of adverse conditions. Plant growth promoting activities are helpful to the plants for their growth and survival in salinity stress condition.

Salinity stress tolerance ability

PG-63 bacterial isolate has grown well in nutrient medium with 1.0 to 15.0% (w/v) NaCl (Table-2). It was observed that the number of bacteria decreased with increased NaCl concentrations, but it could survive at higher NaCl concentrations. PG-63 show the best growth (about 50%) up to 5% NaCl concentration. (37) observed that panchagavya isolates, *Bacillus tequilensis*, and *Bacillus aryabhatai* can be employed in higher saline soils due to their capacity to survive at higher salt concentrations (12% w/v).

Plant growth promoting traits

PG-63 was screened for various plant growth promoting activities like IAA production, GA production, NH₃ production, phosphate solubilization in the presence of NaCl concentration. All the plant growth promoting activities were done with 2%w/v NaCl concentration and 1%v/v inoculum in PG-63.

- **Qualitative estimation of plant growth promoting traits**

Qualitative estimation was done for the primary screening of plant growth-promoting traits (Figure-1). A pink colour was observed, which indicates positive IAA production. The wine-red colour indicated positive GA production. Positive ammonia production indicated a dark yellow colour. The blue colour indicated a positive result for phosphate solubilization.

IAA production

Figure-2 shows IAA production of PG-63 at different time intervals with 2%w/v NaCl concentration. In abiotic stress conditions, PGPB produces an appropriate amount of IAA hormone, which affects root development and growth.

PG-63 shows higher production (22.67 µg/ml) after 48 hours in control, while it shows higher production (25.15 µg/ml) after 120 hours in the presence of 2% NaCl concentration. *JPVS11* strain shows IAA activity even in the presence of high salt concentration (22). (7) reported two PGPB isolates producing 12.0 µg/ml and 7.0 µg/ml IAA after 72 hours. (3) also reported a rhizospheric bacteria that was able to produce 55.90 µg/ml IAA. Phytohormonal analysis of *AF7* revealed 36.41 µg/ml IAA production at 2.5%(w/v) NaCl (10).

- **GA production**

Figure-3 shows GA production of PG-63 different time intervals with NaCl concentration, respectively. GA has a crucial role in cell growth and division, root, stem, and leaf expansion, seed germination or dormancy, fruit senescence and flowering.

PG-63 shows higher production (3.38 mg/ml) in control and 2.19 mg/ml in the presence of NaCl after 48 hours. (10) observed GA production in the *AF7* strain (*Alcaligenes* sp.) in the presence of NaCl concentration. (36) reported two bacterial isolates from the rhizosphere which produced 0.48 mg/mL and 0.419 mg/mL gibberellic acid after 72 hours. *Pseudomonas stutzeri* isolated from rhizospheric soil produced 0.03 mg/ml GA after 72 h (28).

- **NH₃ production**

Figure-4 shows ammonia production of PG-63 at different time intervals in the presence of NaCl concentration. Abiotic stress may restrict the ability of plants to reduce and assimilate nitrogen through the inhibition of enzymes implicated in nitrogen metabolism, such as Nitrite Reductase. Use of PGPB that produces ammonia may provide a biological alternative for fixing atmospheric N₂.

PG-63 shows higher production (36.0 µmol/ml) in control and 33.8 µmol/ml in the presence of NaCl after 48 hours. Bacterial strains *P. reactans* and *P. alli* show ammonia production in the presence of different NaCl concentrations (27). *Azotobacter* isolates obtained from commercial biofertilizers reported 4.75 µg/mL ammonia production (38). *Bacillus megaterium* isolated from soil produced 2.04 µmol/ml ammonia after 96 h (18). *Klebsiella pneumoniae* from chickpea rhizosphere 1.00 µmol/ml ammonia after 96 h (24).

- **Phosphate solubilization**

Figure-5 shows phosphate solubilization of PG-63 at different time intervals in the presence of NaCl concentration. The phosphate-solubilizing microorganisms (PSMs) such as phosphate-solubilizing bacteria (PSB) present in most soils which can solubilize insoluble forms of phosphates.

PG-63 shows higher production (64.40 µg/ml) in control after 10 days, while it shows higher production (70.67 µg/ml) in the presence of NaCl after 3 days. BAM-4, BAM-6, BAM-12, and BAM-77 (soil bacteria) show phosphate solubilization at different NaCl concentrations (15). The *Bacillus cereus* was able to solubilize 159 µg/ml tri-calcium phosphate after 10 days in Pikovskaya's broth (34).

Seed germination test

Table-3 shows seed germination percentage, seedling length (cm) and seed vigour index of *Trigonella foenum-graecum*. One of the most crucial stages of the plant growth cycle is seed germination.

The percentage germination ability of PG-63 was shown in figure-6. *Trigonella foenum-graecum bicolor* seeds were treated with distilled water and showed 82% seed germination with 106.6 seed vigor index. Seeds treated with NaCl showed 45% seed germination with 4.5 seed vigor index. When seeds treated with NaCl along with PG-63 bacterial culture it showed 90% seed germination with 81 seed vigor index. Alfalfa (*Medicago sativa*) showed better seed germination after application of PGPB (*Rhizobia strain*) in salinity stress (2).

Pot study on *Trigonella foenum-graecum*

Table-4 shows physico-vegetative parameters of *Trigonella foenum-graecum*. The biological contribution of PG-63 on the growth of *Trigonella foenum-graecum* was investigated in the presence of NaCl concentration.

The results were recorded with positive and negative control. An increase in root length and shoot length was observed on 15th day along with increase in leaf length (figure-7). It was observed that treated plants had longer roots (12%) and shoots (12.5%) than the untreated control plants (Figure-10). In addition, treated plants had a higher fresh weight and dry weight.

Arachis hypogea that has been exposed to *Bacillus sp. PG-8* showed significant plant growth. A potential method for increasing plant growth and decreasing the use of hazardous chemical fertilisers is to inoculate soil or crops with PGPB. *Arachis hypogea* may have better growth characteristics because of *Bacillus sp. PG-8* in the production of auxins, gibberellic acid, and phosphate solubilization, which increases the availability of nutrients (13).

Table-1: Pot study design for *Trigonella foenum-graecum*

Plant	Pot number	Pot design
<i>Trigonella foenum-graecum</i>	Pot-1	Only seeds
	Pot-2	Seeds+NaCl
	Pot-3	Seeds+PG-63
	Pot-4	Seeds+NaCl+PG-63

Table-2: The percentage growth of PG-63 in salinity stress condition

NaCl (%W/V)	Number of cells	Percent growth (%)
Control	94.67×10 ⁸	100
1%	87.29×10 ⁸	92.2
3%	56.51×10 ⁸	59.7
5%	48.50×10 ⁸	51.2
7%	31.31×10 ⁸	33.0
10%	2.400×10 ⁸	2.53
15%	0.150×10 ⁸	0.12

Table-3: Seed germination and seed vigour index

Treatment	Number of seeds sown	Number of germinated seeds	Germination %	Seedling length (cm)	Vigour index
Only seeds (distilled water)	100	82	82	1.3	106.6
Seeds + 2% NaCl	100	45	45	0.1	4.500
Seeds+2% NaCl + PG-63	100	90	90	0.9	81.00

Table-4: Physico-vegetative parameters of *Trigonella foenum-graecum*

Plant parameters of <i>Trigonella foenum-graecum</i> after 15 days	Control	NaCl (Negative control)	PG-63 (Positive Control)	NaCl+PG-63
Shoot length (cm)	4.000	2.900	4.500	3.000
Root length (cm)	7.000	5.450	7.900	5.570
Leaf length (cm)	3.230	2.430	3.550	2.430
Fresh weight (gm)	0.200	0.120	0.165	0.134
Dry weight (gm)	0.030	0.010	0.025	0.018




Figure-1: Qualitative estimation of plant growth promoting traits along with blank a) IAA production b) GA production c) Ammonia production d) Phosphate solubilization




Figure-2: IAA production by PG-63 in salinity stress along with control




Figure-3: GA production by PG-63 in salinity stress along with control




Figure-4: Ammonia production by PG-63 in salinity stress along with control




Figure-5: Phosphate solubilization by PG-63 in salinity stress along with control




Figure-6: *Trigonella foenum-graecum* seeds a) Only seeds (distilled water) b) seeds+NaCl+PG-63




Figure-7: *Trigonella foenum-graecum* plant after 15 days a) control (only seeds) b) positive control (seeds+PG-63) c) negative control (seeds+NaCl) d) seeds+NaCl+PG-63

CONCLUSION

The current study indicates the importance of PG-63 isolated from fermented panchagavya. PG-63 helps plants to survive in adverse condition like salinity stress because it shows many plant growth promoting activities (IAA production, GA production, ammonia production, phosphate solubilization etc.) even in the salinity stress. The application of PG-63 on plants improved the growth with NaCl concentration. This study can be further explored by application of PG-63 in the field study. PG-63 may be used as biofertilizer. PG-63 isolate may have the potential to re-fertile the saline and sodic soils.

Acknowledgement

The authors duly acknowledge the Department of Microbiology and Biotechnology, Gujarat University, Ahmedabad, Gujarat for providing the laboratory and infrastructure facilities.

Competing interests

The authors have declared that no competing interest exists.

REFERENCES

1. Barnawal, D., Bharti, N., Maji, D., Chanotiya, C. S., & Kalra, A. (2014). ACC deaminase-containing *Arthrobacter protophormiae* induces NaCl stress tolerance through reduced ACC oxidase activity and ethylene production resulting in improved nodulation and mycorrhization in *Pisum sativum*. *Journal of plant physiology*, 171(11), 884-894.
2. Bertrand, A., Dhont, C., Bipfubusa, M., Chalifour, F. P., Drouin, P., & Beauchamp, C. J. (2015). Improving salt stress responses of the symbiosis in alfalfa using salt-tolerant cultivar and rhizobial strain. *Applied Soil Ecology*, 87, 108-117.
3. Bhandari, H., & Khunt, M. D. (2019). Screening of Multi-trait PGPR from Rhizosphere of Sugarcane (*Saccharum officinarum* L.). *Int. J. Curr. Microbiol. App. Sci*, 8(6), 617-623.
4. Bhise, K. K., Bhagwat, P. K., & Dandge, P. B. (2017). Synergistic effect of *Chryseobacterium gleum* sp. SUK with ACC deaminase activity in alleviation of salt stress and plant growth promotion in *Triticum aestivum* L. *3 Biotech*, 7, 1-13.

5. Bric, J. M., Bostock, R. M., & Silverstone, S. E. (1991). Rapid in situ assay for indoleacetic acid production by bacteria immobilized on a nitrocellulose membrane. *Applied and environmental Microbiology*, 57(2), 535-538.
6. Cappuccino, J. G., & Sherman, N. (1992). Biochemical activities of microorganisms. *Microbiology, A Laboratory Manual. The Benjamin/Cummings Publishing Co. California, USA*, 76.
7. Das, S., Nurunnabi, T. R., Parveen, R., Mou, A. N., Islam, M. E., Islam, K. M. D., & Rahman, S. M. (2019). Isolation and characterization of indole acetic acid producing bacteria from rhizosphere soil and their effect on seed germination. *International Journal of Current Microbiology Applications in Science*, 8(3), 1237-1245.
8. Egamberdieva, D., Wirth, S., Bellingrath-Kimura, S. D., Mishra, J., & Arora, N. K. (2019). Salt-tolerant plant growth promoting rhizobacteria for enhancing crop productivity of saline soils. *Frontiers in microbiology*, 10, 2791.
9. EL Sabagh, A., Hossain, A., Barutçular, C., Iqbal, M. A., Islam, M. S., Fahad, S., ... & Erman, M. (2020). Consequences of salinity stress on the quality of crops and its mitigation strategies for sustainable crop production: an outlook of arid and semi-arid regions. *Environment, climate, plant and vegetation growth*, 503-533.
10. Fatima, T., Mishra, I., Verma, R., & Arora, N. K. (2020). Mechanisms of halotolerant plant growth promoting *Alcaligenes sp.* involved in salt tolerance and enhancement of the growth of rice under salinity stress. *3 Biotech*, 10, 1-12.
11. Graham, H. D., & Thomas, L. B. (1961). Rapid, simple colorimetric method for the determination of micro quantities of gibberellic acid. *Journal of Pharmaceutical Sciences*, 50(1), 44-48.
12. Gohil, R. B., Patel, K., Shah H., Vyas, S., Desai, M., Rajput, K., (2020). Isolation and Screening of Plant Growth Promoting Bacteria from Fermented Panchagavya. *Biosc.Biotech.Res.Comm*, 13(1), 42-48.
13. Gohil, R. B., Raval, V. H., Panchal, R. R., & Rajput, K. N. (2022). Plant growth-promoting activity of *Bacillus sp.* PG-8 isolated from fermented panchagavya and its effect on the growth of *arachis hypogea*. *Frontiers in Agronomy*, 4, 11.
14. Herger, G., Nielsen, R., & Margheim, J. (2015). Fertilizer history P3: in WWII nitrogen production issues in age of modern fertilizers.
15. Jha, A., Saxena, J., & Sharma, V. (2013). Investigation on phosphate solubilization potential of agricultural soil bacteria as affected by different phosphorus sources, temperature, salt, and pH. *Communications in soil science and plant analysis*, 44(16), 2443-2458.
16. Jha, Y., & Subramanian, R. B. (2014). PGPR regulate caspase-like activity, programmed cell death, and antioxidant enzyme activity in paddy under salinity. *Physiology and Molecular Biology of Plants*, 20, 201-207.
17. Kalaji, H. M., Bosa, K., Kościelniak, J., & Żuk-Gołaszewska, K. (2011). Effects of salt stress on photosystem II efficiency and CO₂ assimilation of two Syrian barley landraces. *Environmental and Experimental Botany*, 73, 64-72.
18. Krunal, M., & Sanjay, J. (2020). Bacilli consortia positively regulates agronomic and growth traits in rice. *International Journal of Chemical Studies*, 8(3), 636-645.
19. Kumar, A., & Verma, J. P. (2018). Does plant—microbe interaction confer stress tolerance in plants: a review? *Microbiological research*, 207, 41-52.
20. Kumar, A., Singh, S., Gaurav, A. K., Srivastava, S., & Verma, J. P. (2020). Plant growth-promoting bacteria: biological tools for the mitigation of salinity stress in plants. *Frontiers in Microbiology*, 11, 1216.
21. Kumar, K., Verma, G., Veer, R., Kumar, S., & Kumar, P. (2020). Exploitation of Panchagavya, benefits and eco-friendly management of plant diseases: A review. *J. Entomol. Zool. Stud*, 8, 2360-2364.
22. Kumar, A., Singh, S., Mukherjee, A., Rastogi, R. P., & Verma, J. P. (2021). Salt-tolerant plant growth-promoting *Bacillus pumilus* strain JPVS11 to enhance plant growth attributes of rice and improve soil health under salinity stress. *Microbiological Research*, 242, 126616.
23. Lambers, H. (2003). Introduction: Dryland salinity: A key environmental issue in southern Australia. *Plant and soil*, v-vii.
24. Mazumdar, D., Saha, S. P., & Ghosh, S. (2018). Klebsiella pneumoniae rs26 as a potent PGPR isolated from chickpea (*Cicer arietinum*) rhizosphere. *Pharm. Innovat. Int. J*, 7, 56-62.
25. Mehta, P., Jajoo, A., Mathur, S., & Bharti, S. (2010). Chlorophyll a fluorescence study revealing effects of high salt stress on Photosystem II in wheat leaves. *Plant physiology and biochemistry*, 48(1), 16-20.
26. Miceli, A., Vetrano, F., Sabatino, L., D'Anna, F., & Moncada, A. (2019). Influence of preharvest gibberellic acid treatments on postharvest quality of minimally processed leaf lettuce and rocket. *Horticulturae*, 5(3), 63.
27. Moreira, H., Pereira, S. I., Vega, A., Castro, P. M., & Marques, A. P. (2020). Synergistic effects of arbuscular mycorrhizal fungi and plant growth-promoting bacteria benefit maize growth under increasing soil salinity. *Journal of Environmental Management*, 257, 109982.
28. Patel, T., & Saraf, M. (2017). Biosynthesis of phytohormones from novel rhizobacterial isolates and there in vitro plant growth-promoting efficacy. *Journal of Plant Interactions*, 12(1), 480-487.
29. Pikovskaya, R. I. (1948). Mobilization of phosphorus in soil in connection with vital activity of some microbial species. *Mikrobiologiya*, 17, 362-370.
30. Ramadoss, D., Lakkineni, V. K., Bose, P., Ali, S., & Annapurna, K. (2013). Mitigation of salt stress in wheat seedlings by halotolerant bacteria isolated from saline habitats. *SpringerPlus*, 2(1), 1-7.
31. Ram, A. A. M. (2017). Panchagavya is a bio-fertilizer in organic farming. *Int J Adv Sci Res*, 2(5), 54-57.
32. Rütting, T., Aronsson, H., & Delin, S. (2018). Efficient use of nitrogen in agriculture. *Nutrient Cycling in Agroecosystems*, 110, 1-5.
33. Sarkar, A., Pramanik, K., Mitra, S., Soren, T., & Maiti, T. K. (2018). Enhancement of growth and salt tolerance of rice seedlings by ACC deaminase-producing *Burkholderia sp.* MTCC 12259. *Journal of plant physiology*, 231, 434-442.

34. Salwan, R., Sharma, V., Saini, R., & Pandey, M. (2021). Identification of plant beneficial *Bacillus spp.* for Resilient agricultural ecosystem. *Current Research in Microbial Sciences*, 2, 100046.
35. Sharma, S., Kulkarni, J., & Jha, B. (2016). Halotolerant rhizobacteria promote growth and enhance salinity tolerance in peanut. *Frontiers in microbiology*, 7, 1600.
36. Sharma, S., Sharma, A., & Kaur, M. (2018). Extraction and evaluation of gibberellic acid from *Pseudomonas sp.*: Plant growth promoting rhizobacteria. *Journal of Pharmacognosy and Phytochemistry*, 7(1), 2790-2795.
37. Shultana, R., Kee Zuan, A. T., Yusop, M. R., Saud, H. M., & El-Shehawi, A. M. (2021). *Bacillus tequilensis* strain 'UPMRB9' improves biochemical attributes and nutrient accumulation in different rice varieties under salinity stress. *PLoS One*, 16(12), e0260869.
38. Sumbul, A., Ansari, R. A., Rizvi, R., & Mahmood, I. (2020). Azotobacter: A potential bio-fertilizer for soil and plant health management. *Saudi journal of biological sciences*, 27(12), 3634-3640.

Copyright: © 2021 Society of Education. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.