

REVIEW ARTICLE

Regioselective drug delivery system of suitable drug for sustained therapeutic effect by novel drug release mechanism

¹ Mlind Dilip Kalaskar *, ¹Lokesh G. Barde, ¹Wasim G. Shaikh, ¹Anis A. Shaikh, ¹Amol D. landge, ²Harshad P. Khade

¹Department of Pharmaceutics, SSBT'S Institute of Pharmacy, Jalgaon, Maharashtra, India

²Department of Pharmaceutical Chemistry, Annasaheb Dange College of B Pharmacy, Ashta, Maharashtra, India

ABSTRACT

The current project is aimed at the development, formulation, and testing of a regioselective drug delivery system which is to provide the achievement of a long-term therapeutic effect by a new drug release mechanism. The traditional drug delivery methods tend to experience low site specificity, high rate of drug release, and low therapeutic efficacy which contributes to high dosage, and side effects. To address these shortcomings, a specialized delivery methodology has been established based on the high efficiency of carrier systems able to regulate the drug delivery to the specific location of action. The compound of choice was placed in a customized delivery system to increase localization, augment bioavailability and longevity of action. Optimization of the formulation was carried out based on several physicochemical parameters that included particle size, encapsulation percentage, drug content, and in vitro release properties. The system created revealed the controlled and sustained release of drugs which proved the successfulness of the new release mechanism. The studies carried out on evaluation showed that there was an improvement in stability, increased regioselective targeting, and extended therapeutic performance with respect to the traditional formulations. The data indicate that the regioselective drug delivery system developed can provide a promising solution in the treatment of chronic diseases with the lower frequency of doses and the minimization of the side effects of the system. This new method has high potential in enhancing therapeutic results and adherence of patients to targeted therapy of drugs.

Keywords; Regioselective drug delivery; Targeted drug delivery systems; Sustained release formulations; Polymeric nanoparticles; Bio-responsive carriers.

Received 10.02.2026

Revised 21.02.2026

Accepted 20.03.2026

How to cite this article:

Mlind Dilip K, Lokesh G. B, Wasim G. S, Anis A. S, Amol D. I, Harshad P. K. Regioselective drug delivery system of suitable drug for sustained therapeutic effect by novel drug release mechanism . Adv. Biores., Vol 17 (3) March 2026: 64-73.

INTRODUCTION

The establishment of efficient drug delivery systems is becoming more and more popular in modern pharmaceutical research in order to increase the therapeutic effect and reduce the side effects. The traditional dosage forms tend to deliver drugs uncontrollably, thus causing variations in drug concentration, inefficiency and non-compliance of the patient in relation to drug dosing because of frequent dosages. These restrictions have stimulated the development of new delivery methods that can have the capacity to retain the concentration of drugs in the therapeutic range over long intervals. Advanced delivery systems can enhance predictable and effective therapeutic outcomes through enhancement of drug stability, absorption and bioavailability [1].

In order to counter these setbacks, there has been development of controlled and targeted drug delivery strategies which promise to counter the challenges. Such systems allow the site-specific drug release and the duration of therapeutic activity to be lowered, decreasing systemic levels of side effects and enhancing patient compliance. Regioselective delivery gives a high level of localization of the drugs to the target tissues and sustained release formulations extend the availability of a drug and reduces the frequency of drug dosing. Collectively, these measures symbolize a significant breakthrough in

pharmaceutical technology, safer and more effective treatment methods that bring more effective and patient-friendly treatment to a large variety of medical disorders [2].

Introduction to Targeted Drug Delivery.

Targeted drug delivery is described as the process of delivering therapeutic agents to the site of activity with minimal exposure to the non-target tissues. This mechanism increases the amount of drug that is required at the specific site, increases bioavailability and minimizes systemic side effects. There are different carrier-based systems including liposomes, nanoparticles, microspheres and vesicular systems, which have been designed to deliver in a targeted fashion. The advanced platforms can enable a controlled transport, prevent drug degradation and enhance cellular uptake, which makes them very useful in contemporary pharmacotherapy [3].

Drawbacks of traditional methods of drug delivery

The traditional modes of drug delivery such as tablets, capsules and injections usually disseminate drugs in the body without being specific. This is non-selective distribution which might cause low drug concentration at the target site, high frequency of dosing, and adverse effects. Also, quick elimination and metabolism of drugs decrease their therapeutic benefits. These restrictions have led to the necessity of novel drug delivery systems that could offer controlled release, enhanced targeting and sustained drug action [4].

Idea of Regioselective Drug Delivery.

The concept of regioselective drug delivery applies in situations where a specific part or tissue of the body is targeted by the drug to achieve the greatest therapeutic effect and the least possible side effects. It is based on the specific carriers and release mechanisms that react to the physiological conditions of pH, enzymes, temperature or contact with receptors. Having the ability to target the drug to the affected area, regioselective systems benefit the treatment efficacy, decreases the toxicity, and ensures patient compliance which also is a significant breakthrough in the pharmaceutical science [5].

Noteworthiness of Sustained Therapeutic Systems.

Sustained release drug delivery systems help in releasing the drug at a slow rate to achieve a constant concentration of the drug in the body. This limits the variation of the peaks and troughs, lowers dose frequency and patient compliance with treatment. The long-term therapeutic systems are especially useful in chronic diseases that need long-term medication since they enhance the effectiveness of the therapeutic process and lower the number of side effects and the total cost of the treatment [6].

Scope and Significance of Novel Release Mechanisms

New release mechanisms of drugs are crucial in the regulation of drug release rate, location and duration. Such mechanisms can include stimuli-responsive systems, biodegradable polymers, vesicular carriers or smart materials that are responsive to physiological stimuli. The innovation of these highly sophisticated systems provides better stability of the drug, accurate targeting and better therapeutic results. Regioselective targeting combined with sustained release is an innovative approach to drug delivery systems of next generation that has broad potential in pharmaceutical and clinical applications [7].

Regioselective Drug delivery Principles.

Regioselective drug delivery is a new pharmaceutical approach that is designed to deliver therapeutic agents to particular locations in the body with the aim of having optimum therapeutic effect and minimal systemic exposure. This method combines the principles of pharmacokinetics, biological interactions as well as controlled release technologies to overcome the physiological barriers and achieve a certain localization of drug action [8].

Definition and Concept

The process of targeted transport and controlled delivery of drugs to a specific anatomical or physiological location in the body is known as regioselective drug delivery. In contrast to traditional systems where drugs are distributed in an identical manner, regioselective systems are developed to localize the drug in the targeted area with the help of specific carriers, stimuli-responsive materials, or molecular recognition, etc. Such selective localization increases therapeutic effects, decreases dose levels and minimizes adverse effects [9].

Barriers of the Biology and Targeting Problems.

The human body has various biological barriers which limit movements and absorption of the drug and these include the skin, gastrointestinal mucosa, blood-brain barrier, cellular membranes, and enzymatic systems. These obstacles tend to inhibit the drug penetration and lower bioavailability at the target site. Also, targeted delivery is further complicated by non-specific distribution, high metabolism, and immune system elimination. These obstacles need to be overcome by developing novel strategies of formulation like permeation enhancers, vesicular system, biodegradable carriers and any of the receptor-based targeting mechanisms [10].

Pharmacokinetic and Pharmacodynamic.

To a large extent, pharmacokinetic aspects like absorption, distribution, metabolism and excretion contribute to the effectiveness of regioselective drug delivery systems. An effective system must maximize the concentration of drugs at the site of action and have minimal systemic exposure. The pharmacodynamic aspect includes the possibility to obtain the intended therapeutic effect by using reduced doses of the drugs due to an increase in localization. Controlled release kinetics, drug stability, and carrier-drug interactions are of significance in ensuring that there is a longer therapeutic level and better clinical outcomes [11].

Site-Specific Systems vs Regioselective Systems.

Site-specific drug delivery is defined as the specific delivery of drugs to very localized organs, tissues or cells including tumor tissues or targeted receptors, frequently via ligand-mediated processes. Comparatively, regioselective drug delivery is aimed at the delivery of drugs to a more widespread part of anatomical location, e.g. gastrointestinal tract or skin layers or localized inflammatory areas. Although both methods are intended to facilitate an increase in the efficiency of targeting, regioselective systems focus on localization in a region instead of single cellular target, thus being especially efficient in the case of sustained and localized therapeutics [12].

Guidelines to the Choice of Appropriate Drug Candidates.

The effectiveness of a regioselective and sustained drug delivery system mainly relies on the good choice of drug candidates. The targeted and controlled release drugs should have certain physicochemical, biopharmaceutical, and stability properties, which guarantee efficient formulation, therapeutic functionality, and patient safety. Close consideration of these attributes promotes drug loading performance, predictable release pattern, and the system stability [13].

Physicochemical Properties

The physicochemical characteristics of molecular weight, lipophilicity, melting point, and ionization constant have a great impact on the drug permeability, encapsulation efficiency, and release profile. Medium-weight drugs that have a balanced hydrophilic-lipophilic characteristic are the favored drugs to be used in regioselective delivery systems. These attributes promote breaching biological obstacles and fixed assimilation into carrier systems, which allow regulated and consistent drug delivery [14].

Biopharmaceutical Considerations

Such biopharmaceutical parameters as absorption rate, bioavailability, metabolism as well as elimination half-life are important in determining the suitability of the drug. Sustained and targeted delivery systems are advantageous to drugs that have a low bioavailability, high metabolism, or low half-lives. The alternate method of regioselectivity assists in increasing concentrations of therapy at the point of attack and reducing the systemic clearance, thus increasing the overall effectiveness of therapy [15].

Dose and Therapeutic Window

To permit effective concentrations of drug without resulting in toxicity, an ideal drug candidate must have a clear therapeutic window. Sustained release systems are particularly appropriate to drugs that have to be taken regularly or have a thin range of therapeutic levels. The administration of drugs in a controlled manner serves to regulate plasma levels within the therapeutic range during prolonged intervals reducing variations and decreasing the possibility of adverse effects [16].

Solvability Factor and Stability Factor.

The stability of drugs during formulation and physiological processes is what is needed to preserve therapeutic effects. The drug must be chemically and physically stable through processing, storage and release. Solubility is also known to affect the drug loading and kinetics of drug release; ill soluble drugs can be subject to the formulation strategy of solubilizing and/or encapsulating it in carriers to boost bioavailability and regulated drug release behavior [17].

Drug-Carrier Compatibility

The interaction of the drug and the carrier system of choice is critical in order to achieve stability in formulation and appropriate release properties. There should be no interactions that lead to the degradation, precipitation or loss of activity of the drug. Compatibility experiments based on methods like the difference scanning calorimetry, Fourier-transform infrared spectroscopy and stability analysis are applicable to ensure that the drug is not degraded and it is uniformly spread in the delivery system [18].

New Drug Release Profiles.

The concept of controlled, sustained and regioselective therapeutic effects of advance drug delivery is attained by the effect of novel drug release mechanisms. These processes focus on the rate, duration and the site of drug liberation using physical, chemical and biological stimuli. Such systems have been

developed and the stability of drugs, target efficiency and patient compliance have been greatly enhanced [19].

Diffusion-Controlled Systems

Diffusion-controlled drug delivery systems entrap the drug using a polymeric matrix or membrane as the release system is gradual release of the drug depending on the concentration gradients. The drug diffuses slowly out of the carrier to the biological environment to deliver the longest levels of treatment. They are extensively employed because they are simple, have predictable release kinetics and are able to achieve stable drug concentrations with time [20].

Erosion-Controlled Systems

Drug release in systems controlled by erosion is by the gradual degradation or dissolution of the polymeric carrier in physiological conditions. The release profile is defined by the erosion rate, making it possible to release drugs at a constant and even at a zero-order rate. Such systems usually utilize biodegradable polymers, and these polymers are especially applicable to long-term and implantable drug delivery systems [21].

Stimuli-Responsive Systems

Smart drug delivery systems are stimuli-responsive or smart, i.e., drug delivery systems that release drugs based on a particular stimulus (physiological or external) [22].

pH-Sensitive Systems

These systems take advantage of the differences in pH between different parts of the body, including acidic tumor microclimate or the GIT. The controlled release of the drug is caused when the carrier is exposed to a certain pH, which allows the targeting of controlled delivery [23].

Temperature-Sensitive Systems

Temperature sensitive systems make use of polymers that experience a phase transition at certain temperatures and release drugs when subjected to heat, whether in the form of physiological or external heat. These systems can be applicable in localized hyperthermic therapies [24].

Enzyme-Triggered Systems

Enzyme-sensitive carriers are drugs that liberate under the action of certain enzymes that are present at the sites of the disease, i.e. inflamed tissue or tumor. The method varies the drug release to sites and the exposure to the system [25].

Redox-Sensitive Systems

Redox-responsive systems are stimulated by changes in the intracellular and extracellular redox state, especially high levels of glutathione in the cell. With the help of these systems, intracellular drug release can be controlled, which increases the efficacy of therapies [26].

The Osmotically Controlled Release.

The osmotic drug delivery systems rely on the differences in osmotic pressure to maintain the rate of drug release at a constant rate. Water gets into the system via a semi permeable membrane which creates pressure that causes the drug to push out via a delivery orifice. The systems offer very predictable and extended release of drugs which are not very dependent on the conditions of the physique [27].

Smart Polymeric Systems

The Smart polymeric systems are means to combine controlled release, environmental responsiveness and biodegradability. These polymers are able to conform to the physiological conditions thus providing accurate drug delivery over a long time. Their tunable properties make them ideal in regioselective targeting, sustained therapy and improved therapeutic outcomes in current drug therapies [28].

Carrier Systems Used in Regioselective Delivery

Carrier systems are the basis of regioselective drug delivery and they allow the targeted localization of therapeutic agents, regulated release and protect therapeutics. These systems enhance stability of drugs, increases penetration of the drug through the biological barriers and the retention of therapeutic concentrations in particular areas [29].

Synergy in Regioselective Drug Delivery




Figure 1. Synergy in Regioselective Delivery

Polymeric Nanoparticles

Polymeric nanoparticle is a carrier that is biodegradable and can be used to incorporate drugs inside polymer matrix so that they are released in a controlled manner and last longer. They are nanoscale, which facilitates greater tissue penetration and efficient drug delivery to areas where it is required [30].

Lipid-Based Systems

Carriers based on lipids like liposomes and solid lipid nanoparticles increase bio-compatibility and solubility of drugs. They replicate biological membranes, enhance organizational cellular absorption, and targeted therapeutic effect [31]

Vesicular Systems

Vesicular systems are bilayer systems that have the capacity of being loaded with hydrophilic and lipophilic drugs. These systems are able to offer better permeability, release control, and localization of these systems in specific tissues [32].

Hydrogels and In Situ Gels

Hydrogels are three dimensional networks that entrap drugs and release them over time whereas in situ gels release through a conversion to gel form in physiological conditions. These systems provide long retention of drug at the point of delivery [33].

Microspheres and Implants

Microspheres allow long term diffusion of drugs or erosion of polymer, whereas implants allow long term treatment during weeks or months. They are particularly applied to chronic treatment [34].

Hybrid Nanocarriers

Hybrid nanocarriers refer to polymeric and lipid-based nanocarriers that are developed to achieve a high stability, targeting and controlled release. Such multifunctional systems are one of the novel platforms of selective therapy [35].

Formulation Strategies

Regioselective and sustained drug delivery systems require formulation strategies to develop effective systems. A step-by-step systematic and scientific methodology is used to guarantee optimum loading of the drug, a regulated release profile, stability, and targeting. Current development of formulations combines selection of material, carrier engineering and statistics optimization to obtain high efficacy [36].

Rational Design Approach

The rational approach to design aims at attaining knowledge on drug properties, biological barriers and carrier behavior prior to formulation development. With the help of the analysis of physicochemical properties and therapeutic goals, appropriate delivery systems and release mechanisms are chosen. Such an approach minimizes the trial-and-error exploration and enhances the performance of formulation and reproducibility [38].

Selection Criteria of Excipients.

Excipients are selected according to biocompatibility, stability, acceptance by regulatory authorities and as substances that will increase solubility, permeability and release inhibition. Commonly used to enhance formulation performance is the use of surfactants, stabilizers, polymers as well as penetration enhancers to ensure safety and compatibility [38].

Selection and Functionalization of Polymers.

Polymers are involved in regioselective and sustained drug delivery. The process is based on biodegradability, mechanical strength and release properties. Controlling and targeted drug release in response to physiological conditions Polymers targeted Functionalization of polymers with responsive groups or surface modifiers enhances targeting and controlled drug release [39].

Targeting Ligand Incorporation.

Locating ligands like peptides, antibodies or receptor-specific molecules are attached on carrier surfaces to enhance site-specific localization. These ligands improve cellular specificity and cellular uptake and hence promote improved therapeutic efficacy and minimization of systemic exposure [40].

Design of Experiments (DoE) Optimization.

DoE is a statistical method that seeks to analyze the input variables in formulation on key quality parameters like particle size, drug loading and release profile. It facilitates orderly optimization by doing as few experiments as possible to establish strong and repeatable formulation development [41].

Methods of Preparation

The preparation methodology has a significant impact on physicochemical properties, drug loading ability as well as release behavior of regioselective drug delivery systems. The choice of a suitable fabrication process is made depending on the type of the drug, carrier substance, and the required therapeutic outcome. The established as well as the innovative methods of preparation are used to obtain controlled and reproducible formulations [42].

Evaporation of Solvents Methods.

The process of solvent evaporation entails dissolution of the drug, and polymer in an organic solvent, and emulsification in a water phase. Solvent evaporation causes the formation of solid particles which contain the drug. The approach is common to creating nanoparticles and microspheres that have sustained release characteristics [43].

Emulsification Methods

The emulsification methods produce dispersed particles of one phase in another immiscible phase through the surfactants. The single and dual emulsion systems are used depending on the solubility of the drug. Using these methods, it is possible to efficiently encapsulate both hydrophilic and lipophilic drugs and control the formation of particles [44].

Nanoprecipitation

Nanoprecipitation is an easy and fast technique as a polymer solution of a drug in a non-solvent is applied and the rapid solvent displacement generates the particles immediately. It yields homogenous nanoparticles with a high encapsulation capacity and few processing procedures [45].

Spray Drying

Spray drying is used to turn liquid formulations into dry particulate systems by selectively swiftly drying the solvents in a hot chamber. The approach is a scalable synthesis of controlled release particles of known size which, in turn, can be used in a range of drug delivery projects [46].

Microfluidic Techniques

Microfluidic systems exploit the use of controlled microscale channels to mix formulation components with precision leading to uniform and reproducible nanoparticles. The methods provide a better control of particle size distribution and formulation consistency [47].

The Cutting-edge Fabrication Technologies.

More sophisticated technologies like electrospinning and 3D printing provide the opportunity to produce tailored drug delivery systems with good control of drug distribution and release pattern. Such technologies allow building incredibly specific and effective regioselective delivery platforms [48]

Mechanistic Impressions of Intelligent Drug Given Discharges.

Sustained release systems and regioselective approaches to drug delivery require the mechanistic knowledge of the targeted drug delivery. Such processes regulate the drug carrier passage across biological barriers, cell entry, localization of the drug carrier at the target site, and controlled release of the drugs. The understanding of such processes assists in optimization of formulation and predictive therapeutic outcomes [49].

Transport Mechanisms

Transport mechanisms refer to the way in which drugs or carriers will pass through biological membranes and tissues. They are passive diffusion, facilitated transport, active transport, and paracellular pathways. Advanced carrier systems are used to improve permeation through stability of drugs, enhanced interactions with surfaces and temporary alterations of barrier permeability to enhance regioselective targeting [50].

Cellular Uptake Pathways

Endocytic cellular internalization mainly takes place via a clathrin-mediated uptake, caveolae-mediated transport, macropinocytosis, and receptor-mediated internalization. Selective uptake is augmented by the incorporation of targeting ligands on carrier surfaces that are incorporated selectively in particular cells. Upon internalization, intracellular conditions cause drugs to be released leading to a better therapy [52].

Localization and Retention of Tissues.

Tissue retention The capacity of the drug carriers to stay longer in the targeted part is known as tissue retention. This could be by passive accretion in permeable tissues, specific binding to receptors or by means of interaction between the local extracellular matrices. The extended localization is a guarantee of the long-term therapeutic effect and less systemic exposure [52].

Release Behavior Mathematic Modeling.

Kinetics of drug release out of the delivery systems are mathematically modeled and predicted. The most common ones are zero-order, first-order, diffusion-based and empirical equations representing release mechanisms. These models can be used to gain insights into the release behavior, how to optimize the formulation parameters and to ensure that there is a consistent performance in the therapy [53].

New Technologies and Recent Improvements.

The recent advancements in pharmaceutical sciences have greatly increased the effectiveness and accuracy of the regioselective drug delivery systems. Material science, Biotechnology and computational tools have facilitated the creation of sophisticated carriers which can provide responsive, controlled, and patient-specific therapeutic interventions [54].

Smart Nanocarriers

Smart nanocarriers are designed in a way that they respond to a particular physiological stimulus like pH or temperature or enzyme or redox. These systems stay constant in normal condition and deliver drugs specifically to target site which enhances therapeutic performance and reduces systemic toxicity [55].

Bio-responsive Systems

Bio-responsive delivery systems make use of biological signals that are found in diseased tissues to initiate the release of drugs. They are enzyme sensitive polymers, pH sensitive carriers, and redox sensitive nanoparticles. These systems improve pathological microenvironment exploitation, which improves site-specific targeting and controlled therapeutic action [56].

Individualized Drug Delivery Platforms.

Personalized drug delivery aims at designing the formulations with attention to the specifics of individual patients including genetic profile, disease severity, and pharmacokinetic behavior. Individualized release profiles and targeting plans enhance the outcome of treatments and lessens of adverse reactions, favoring the development of precision medicine [56].

Artificial Intelligence in Formulation Design.

Tools based on artificial intelligence and machine learning are embraced more often to predict performance of formulations, optimize the choice of excipients and model drug release. With these technologies, it takes shorter durations of development, minimizes the cost of conducting experiments and enhances the strength of formulations with respect to making decisions based on data.

Table 1. Recent Innovations in Smart Drug Delivery Technologies and Their Therapeutic Applications

Technology	Key Feature	Mechanism of Action	Major Advantage	Application Area	References
Smart Nanocarriers	Stimuli-responsive materials	Release drug in response to pH, temperature, enzymes, or redox changes	High targeting precision	Cancer therapy, localized infections	[57]
Bio-responsive Systems	Pathology-triggered activation	Drug release in diseased microenvironment	Reduced systemic toxicity	Inflammation, tumors	[58]
Personalized Delivery Platforms	Patient-specific design	Customized dose and release profile	Improved therapeutic outcome	Precision medicine	[59]
Artificial Intelligence (AI)	Data-driven formulation design	Predicts and optimizes formulation parameters	Faster development & accuracy	Drug delivery optimization	[60]

FUTURE PERSPECTIVES

The combination of smart materials, nanotechnology, and biological targeting plans is likely to bring regioselective drug delivery systems to a new level. Bio-responsive carriers that can be used in future to deliver drugs according to certain physiological needs will also enhance more precise therapeutic delivery and minimize systemic side effects. The emergence of personalized medicine will facilitate personalized platforms of delivering treatment as per the needs of the patients so as to improve the result of dose and treatment [61].

In addition, computational modeling and artificial intelligence will be relevant in formulation design and optimization, which will speed up the development and enhance reproducibility. The further development of scalable manufacturing methods including microfluidics and 3D printing will facilitate the commercial fabrication of complicated delivery systems. On the whole, these inventions will put regioselective drug delivery in a more efficient, patient-specific, and clinically effective treatment line [61].

CONCLUSION

Regioselective drug delivery systems can provide a potent method to enhance the level of therapeutic efficacy by releasing drugs to targeted areas and attaining sustained release. These innovative systems are superior to conventional drug dosages, and improve localization of drugs, decrease systemic adverse effects, and elevate patient compliance. Novel release mechanisms, smart polymers and advanced carrier technologies have helped a great deal to reinforce controlled and targeted performance in drug delivery. The aspects of rational formulation strategy, selection of correct drugs, and contemporary preparation methods are important in the provision of stable and efficient delivery platforms. Mechanistic knowledge of the transport routes, the cellular uptake, and release kinetics further assist in optimization of the system and reliable therapeutic results. Regioselective drug delivery is in the process of transforming further to more precise and clinical relevance with the introduction of bio-responsive materials, tailored delivery systems, and the use of artificial intelligence to aid the design of the formulations. On the whole, the systems are a bright future of the safer, more effective, and sustainable therapeutic treatment in various medical applications.

REFERENCES

1. Crommelin, D.J. & Florence, A.T. (2013). Towards more effective advanced drug delivery systems. *Int. J. Pharm.*,454(1):496-511.
2. Bae, Y.H. & Park, K. (2011). Targeted drug delivery to tumors: myths, reality and possibility. *J. Control. Release*,153(3):198-205.
3. Ashique, S., Sandhu, N.K., Chawla, V. & Chawla, P.A. (2021). Targeted drug delivery: trends and perspectives. *Curr. Drug Deliv.*,18(10):1435-1455.
4. Prabahar, K., Alanazi, Z. & Qushawy, M. (2021). Targeted drug delivery system: advantages, carriers and strategies. *Indian J. Pharm. Educ. Res.*,55:346-353.
5. Firer, M.A. & Gellerman, G. (2012). Targeted drug delivery for cancer therapy: the other side of antibodies. *J. Hematol. Oncol.*,5(1):70.
6. Grant, R.L. (1979). The problem-oriented system and record keeping in the behavioral therapies. *J. Community Psychol.*,7(1):53-59.
7. Ding, C. & Li, Z. (2017). A review of drug release mechanisms from nanocarrier systems. *Mater. Sci. Eng. C.*,76:1440-1453.
8. Oh, J.K., Lee, D.I. & Park, J.M. (2009). Biopolymer-based microgels/nanogels for drug delivery applications. *Prog. Polym. Sci.*,34(12):1261-1282.
9. Yamashita, F. & Hashida, M. (2013). Pharmacokinetic considerations for targeted drug delivery. *Adv. Drug Deliv. Rev.*,65(1):139-147.
10. Alonso, M.J. (2004). Nanomedicines for overcoming biological barriers. *Biomed. Pharmacother.*,58(3):168-172.
11. Glassman, P.M. & Muzykantov, V.R. (2019). Pharmacokinetic and pharmacodynamic properties of drug delivery systems. *J. Pharmacol. Exp. Ther.*,370(3):570-580.
12. Vyas, S.P. & Sihorkar, V. (2000). Endogenous carriers and ligands in non-immunogenic site-specific drug delivery. *Adv. Drug Deliv. Rev.*,43(2-3):101-164.
13. Bayliss, M.K., Butler, J., Feldman, P.L., Green, D.V., Leeson, P.D., Palovich, M.R. & Taylor, A.J. (2016). Quality guidelines for oral drug candidates: dose, solubility and lipophilicity. *Drug Discov. Today*,21(10):1719-1727.
14. Edwards, M.P. & Price, D.A. (2010). Role of physicochemical properties and ligand lipophilicity efficiency in addressing drug safety risks. *Annu. Rep. Med. Chem.*,45:380-391.
15. Wagner, J.G. (1961). Biopharmaceutics: absorption aspects. *J. Pharm. Sci.*,50(5):359-387.
16. Paci, A., Veal, G., Bardin, C., Levêque, D., Widmer, N., Beijnen, J., Astier, A. & Chatelut, E. (2014). Review of therapeutic drug monitoring of anticancer drugs part 1—cytotoxics. *Eur. J. Cancer*,50(12):2010-2019.

17. Paci, A., Veal, G., Bardin, C., Levêque, D., Widmer, N., Beijnen, J., Astier, A. & Chatelut, E. (2014). Review of therapeutic drug monitoring of anticancer drugs part 1–cytotoxics. *Eur. J. Cancer*,50(12):2010-2019.
18. Ding, C. & Li, Z. (2017). A review of drug release mechanisms from nanocarrier systems. *Mater. Sci. Eng. C*,76:1440-1453.
19. Varma, M.V., Kaushal, A.M., Garg, A. & Garg, S. (2004). Factors affecting mechanism and kinetics of drug release from matrix-based oral controlled drug delivery systems. *Am. J. Drug Deliv.*,2(1):43-57.
20. Siepmann, J. & Siepmann, F. (2012). Modeling of diffusion-controlled drug delivery. *J. Control. Release*, 161(2):351-362.
21. Zuleger, S. & Lippold, B.C. (2001). Polymer particle erosion controlling drug release. *Int. J. Pharm.*,217(1-2):139-152.
22. Indermun, S., Govender, M., Kumar, P., Choonara, Y.E. & Pillay, V. (2018). Stimuli-responsive polymers as smart drug delivery systems. In: *Stimuli Responsive Polymeric Nanocarriers for Drug Delivery Applications*, Vol.1, Woodhead Publ.,43-58.
23. Zeeshan, M., Ali, H., Khan, S., Khan, S.A. & Weigmann, B. (2019). Advances in orally-delivered pH-sensitive nanocarrier systems. *Int. J. Pharm.*,558:201-214.
24. Choi, S.W., Zhang, Y. & Xia, Y. (2010). A temperature-sensitive drug release system based on phase-change materials. *Angew. Chem. Int. Ed.*,49(43):7904-7908.
25. Khan, M.F., Kundu, D., Gogoi, M., Shrestha, A.K., Karanth, N.G. & Patra, S. (2020). Enzyme-responsive and enzyme immobilized nanoplatfoms for therapeutic delivery. *Nanopharmaceuticals*,3:165-200.
26. Guo, X., Cheng, Y., Zhao, X., Luo, Y., Chen, J. & Yuan, W.E. (2018). Advances in redox-responsive drug delivery systems of tumor microenvironment. *J. Nanobiotechnol.*,16(1):74.
27. Keraliya, R.A., Patel, C., Patel, P., Keraliya, V., Soni, T.G., Patel, R.C. & Patel, M.M. (2012). Osmotic drug delivery system as a part of modified release dosage form. *ISRN Pharm.*,2012:528079.
28. Aghabegi Moghanjoughi, A., Khoshnevis, D. & Zarrabi, A. (2016). A concise review on smart polymers for controlled drug release. *Drug Deliv. Transl. Res.*,6(3):333-340.
29. Ding, C. & Li, Z. (2017). A review of drug release mechanisms from nanocarrier systems. *Mater. Sci. Eng. C*,76:1440-1453.
30. Soppimath, K.S., Aminabhavi, T.M., Kulkarni, A.R. & Rudzinski, W.E. (2001). Biodegradable polymeric nanoparticles as drug delivery devices. *J. Control. Release*,70(1-2):1-20.
31. Patel, D., Solanki, J., Kher, M.M. & Azagury, A. (2024). Surface engineering of lipid-based drug delivery systems. *Small*,20(43):2401990.
32. Jain, S., Jain, V. & Mahajan, S.C. (2014). Lipid based vesicular drug delivery systems. *Adv. Pharm.*,2014:574673.
33. Manoukian, O.S., Arul, M.R., Sardashti, N., Stedman, T., James, R., Rudraiah, S. & Kumbar, S.G. (2018). Biodegradable polymeric injectable implants for long-term delivery of contraceptive drugs. *J. Appl. Polym. Sci.*,135(14):46068.
34. Rizwanullah, M., Alam, M., Harshita, Mir, S.R., Rizvi, M.M. & Amin, S. (2020). Polymer-lipid hybrid nanoparticles: a next-generation nanocarrier for targeted treatment of solid tumors. *Curr. Pharm. Des.*,26(11):1206-1215.
35. Truong-Le, V., Lovalenti, P.M. & Abdul-Fattah, A.M. (2015). Stabilization challenges and formulation strategies associated with oral biologic drug delivery systems. *Adv. Drug Deliv. Rev.*,93:95-108.
36. Sahlgren, C., Meinander, A., Zhang, H., Cheng, F., Preis, M., Xu, C., Salminen, T.A., Toivola, D., Abankwa, D., Rosling, A. & Karaman, D.S. (2017). Tailored approaches in drug development and diagnostics. *Adv. Healthc. Mater.*,6(21):1700258.
37. Elder, D.P., Kuentz, M. & Holm, R. (2016). Pharmaceutical excipients—quality, regulatory and biopharmaceutical considerations. *Eur. J. Pharm. Sci.*,87:88-99.
38. Nicolas, J., Mura, S., Brambilla, D., Mackiewicz, N. & Couvreur, P. (2013). Design, functionalization strategies and biomedical applications of targeted biodegradable polymer-based nanocarriers. *Chem. Soc. Rev.*,42(3):1147-1235.
39. Vyas, S.P. & Sihorkar, V. (2000). Endogenous carriers and ligands in non-immunogenic site-specific drug delivery. *Adv. Drug Deliv. Rev.*,43(2-3):101-164.
40. Beg, S., Swain, S., Rahman, M., Hasnain, M.S. & Imam, S.S. (2019). Application of design of experiments in pharmaceutical product and process optimization. In: *Pharmaceutical Quality by Design*, Academic Press,43-64.
41. Barich, D.H., Munson, E.J. & Zell, M.T. (2005). Physicochemical properties, formulation, and drug delivery. *Drug Deliv. Principles Appl.*,57-71.
42. O'Donnell, P.B. & McGinity, J.W. (1997). Preparation of microspheres by the solvent evaporation technique. *Adv. Drug Deliv. Rev.*,28(1):25-42.
43. Lin, C.Y. & Chen, L.W. (2006). Emulsification characteristics of three- and two-phase emulsions prepared by ultrasonic emulsification method. *Fuel Process. Technol.*,87(4):309-317.
44. Yan, X., Bernard, J. & Ganachaud, F. (2021). Nanoprecipitation as a straightforward process to create polymeric colloidal morphologies. *Adv. Colloid Interface Sci.*,294:102474.
45. Ferreira, M.P., Martins, J.P., Hirvonen, J. & Santos, H.A. (2020). Spray-drying for the formulation of oral drug delivery systems. In: *Nanotechnology for Oral Drug Delivery*, Academic Press,253-284.
46. Yu, M., Mathew, A., Liu, D., Chen, Y., Wu, J., Zhang, Y. & Zhang, N. (2024). Microfluidics for formulation and scale-up production of nanoparticles for biopharma industry. In: *Microfluidics in Pharmaceutical Sciences*, Springer,395-420.

47. Kacem, E., Suk, H.D., Ponnamma, D., Hassan, M.K., Hawari, A., Alshammari, B.A. & Al-Ejji, M. (2025). Exploring 3D printing and electrospinning technologies for advanced porous membrane fabrication. *Adv. Mater. Technol.*,70032.
48. Sharma, R., Singh, D., Gaur, P. & Joshi, D. (2021). Intelligent automated drug administration and therapy: future of healthcare. *Drug Deliv. Transl. Res.*,11(5):1878-1902.
49. Wilson, D.B. (1978). Cellular transport mechanisms. *Annu. Rev. Biochem.*,47(1):933-965.
50. Bishop, N.E. (1997). An update on non-clathrin-coated endocytosis. *Rev. Med. Virol.*,7(4):199-209.
51. Wang, S., Huang, P. & Chen, X. (2016). Hierarchical targeting strategy for enhanced tumor tissue accumulation/retention and cellular internalization. *Adv. Mater.*,28(34):7340-7364.
52. Siepmann, J. & Göpferich, A. (2001). Mathematical modeling of bioerodible polymeric drug delivery systems. *Adv. Drug Deliv. Rev.*,48(2-3):229-247.
53. Crommelin, D.J. & Florence, A.T. (2013). Towards more effective advanced drug delivery systems. *Int. J. Pharm.*,454(1):496-511.
54. Shishir, M.R., Gowd, V., Suo, H., Wang, M., Wang, Q., Chen, F. & Cheng, K.W. (2021). Advances in smart delivery of food bioactive compounds using stimuli-responsive carriers. *Compr. Rev. Food Sci. Food Saf.*,20(6):5449-5488.
55. Patil, A.S., Gadad, A.P. & Dandagi, P.M. (2021). Mono and multi-stimuli responsive polymers: application as intelligent nano-drug delivery systems. *Nanopharm. Adv. Deliv. Syst.*,237-265.
56. Qiao, Y., Wan, J., Zhou, L., Ma, W., Yang, Y., Luo, W., Yu, Z. & Wang, H. (2019). Stimuli-responsive nanotherapeutics for precision drug delivery and cancer therapy. *Wiley Interdiscip. Rev. Nanomed. Nanobiotechnol.*,11(1):e1527.
57. Karati, D. (2023). A concise review on bio-responsive polymers in targeted drug delivery system. *Polym. Bull.*,80(7):7023-7045.
58. Asamoah, A.N., Appiagyei, J.B., Amofa, F.A. & Otu, R.O. (2024). Personalized nanomedicine delivery systems using machine learning and patient-specific data. Syed Khundmir Azmi.
59. Asediya, V., Anjaria, P., Dhial, K. & Pathak, A. (2025). Artificial intelligence and machine learning in drug delivery optimization. In: *Next-Generation Drug Delivery Systems*, Springer,499-521.
60. Summer, M., Hussain, T., Ali, S., Khan, R.R., Muhammad, G. & Liaqat, I. (2025). Exploring the underlying modes of organic nanoparticles in diagnosis, prevention, and treatment of cancer. *Int. J. Polym. Mater. Biomater.*,74(9):829-845.
61. Tiwari, P.C., Pal, R., Chaudhary, M.J. & Nath, R. (2023). Artificial intelligence revolutionizing drug development: exploring opportunities and challenges. *Drug Dev. Res.*,84(8):1652-1663.

Copyright: © 2026 Author. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.