

ORIGINAL ARTICLE

Combining ability and gene action analysis in pearl millet  
[*Pennisetum glaucum* (L) R.Br.]

Reshma Krishnan M. R.<sup>1\*</sup>, M. S. Patel<sup>2</sup> and R. A. Gami<sup>3</sup>

<sup>1</sup>Department of Genetics and Plant Breeding, CPCA, S.D. Agricultural University, Sardarkrushinagar (Gujarat)

<sup>2</sup>Center for crop improvement, S.D. Agricultural University, Sardarkrushinagar (Gujarat)

<sup>3</sup>Maize Research Station, S.D. Agricultural University, Bhiloda-383 245 (Gujarat)

\*Corresponding author **E-mail:** reshmakrishnan086@gmail.com

ABSTRACT

The present investigation was undertaken with the objectives to assess the gene action and magnitude and effects of combining ability of 35 hybrids resulted from Line x Tester mating design with seven lines and five testers and the standard check (GHB 558). The analysis of variance for parents and hybrids was highly significant for all traits indicated the existence of appreciable amount of genetic variability in the experimental material. The magnitude of sca variances was higher than the gca variances for all the characters. This indicated non-additive gene action in the inheritance of these traits. The female ICMA 96222 showed good general combining ability for different characters. The cross ICMA 06777 x 18805 R and ICMA 96222 x 18488 R had good x average combiner parents and good x good combiner parents respectively, high per se performance and positive specific combining ability effect for different characters including grain yield.

**Key Words:** Pear millet, L x T design, combining ability, GCA, SCA.

Received 02.01.2019

Revised 24.01.2019

Accepted 01.03.2019

How to cite this article:

Reshma Krishnan M. R, M. S. Patel and R. A. Gami: Combining ability and gene action analysis in pearl millet [*Pennisetum glaucum* (L) R.Br.]. Adv. Biores. Vol 10 [2] March 2019: 82-86

INTRODUCTION

Pearl millet [*Pennisetum glaucum* (L.) R. Br.] (also known under synonyms: *P. americanum* (L.) Leeke or *P. typhoides* (Burm.) Stapf and C.E. Hubb.) is an important millet crop of traditional farming systems in tropical and subtropical Asia and sub-Saharan Africa. It was introduced to India about 2000 B.C. [1]. Pearl millet is the fourth most important cereal crop in India, after rice, wheat and sorghum, where it is widely grown in the states of Rajasthan, Maharashtra, Gujarat and Haryana where the food security of the poorest population depends vastly on pearl millet production. It contains adequate amount of essential amino acids such as tryptophan, threonine, arginine, and lysine. It is used as a feed, fodder, construction material and even its potential as a source of biofuel is being explored [16].

Discovery of A1 cytoplasmic-nuclear male sterility (CMS) at Tifton, Georgia, USA [4] initiated the era of hybrid cultivar development in pearl millet [*Pennisetum glaucum* (L.) R. Br.], which led to the release of the first grain hybrid in India in 1965 [2]. Since then hundreds of commercial hybrids have been developed and released or commercialized. Hanna [7] identified an A4 CMS system at Tifton, Georgia, USA in a wild grassy *Pennisetum glaucum* (L.) R. Br. subsp. *monodii* (Maire) Brunken. Also, an A5 CMS system was identified in a pearl millet gene pool. Among the various CMS systems reported so far, A4 and A5 CMS systems were found to have the most stable male sterility [14].

Since yield has low heritability, direct selection for yield is not sufficiently effective. So indirect selection for yield has to be done in the light of gene action, by selection of yield contributing traits and with the selection of yield traits. The combining ability studies provide useful information regarding the selection of suitable parents for effective hybridization programme and at the same time elucidates the nature and

magnitude of gene action. In order to proceed with any breeding programme, the breeder has to know about the gene action and the genetic architecture of population. Since, the nature of gene action varies with genetic architecture of population involved in hybridization, it is necessary to evaluate the parents for their combining ability. Evaluation of crosses and identification of the superior cross combinations (combining ability) for yield and quality traits essential for effective manipulation through hybrid breeding.

## MATERIAL AND METHODS

The experimental material consisted of seven maintainer lines used as female lines (ICMA -06777, ICMA -07777, ICMA-96222, ICMA-97111, ICMA-98444, ICMA-04999 and ICMA-05444) and five restorer lines used as testers (J18488-R, 18587-R, 18805-R, 17369-R and 17548-R), crossed in a Line x Tester mating design. The resultant 35 hybrids along with their 12 parents with check GHB-558 were evaluated in Randomized Block Design with three replications at Centre for Crop Improvement, S. D. Agricultural University, Sardarkrushinagar-385 506 (Gujarat) during summer 2016- 2017. The observations were recorded on five randomly selected competitive plants of each genotype in each replication for various characters *i.e.* days to flowering, days to maturity, plant height (cm), number of effective tillers per plant, ear head length(cm), ear head girth(mm), test weight(g), grain yield per plant(g) and harvest index(%). Days to flowering (DF) on the basis of 50% plants of each genotype flowered, days to maturity (DM) on the basis of 80% plants of each genotype matured were recorded. The replication wise mean values were used in statistical analysis. The data were subjected to analysis of variance as per the procedure suggested by Sukhatme and Amble [15]. The combining ability analysis was performed for a Line x Tester mating design as per the method suggested by Kempthorne [9].

## RESULTS AND DISCUSSION

The analysis of variance depicting mean squares due to genotypes were highly significant for all the characters are presented in Table 1. Further, partitioning of the genotypes variance into parents, hybrids and parents vs. hybrids revealed that the parents as well as hybrids exhibited significant differences for all the traits. This indicated the existence of appreciable amount of genetic variability in the experimental material.

Table 1: Analysis of variance (Mean square) for parents and hybrids for seed yield and its component characters in pearl millet

Source of variation	d.f	Days to flowering	Days to maturity	Plant height (cm)	No. of effective tillers per plant	Ear head length (cm)	Ear head girth(mm)	Test weight (g)	Grain yield per plant(g)	Harvest index (%)
Replication	2	3.13	1.68	866.36	0.03	1.65	3.51	0.35	9.53	2.89
Genotype	46	71.08**	86.15**	5163.65**	1.00**	38.68**	72.66**	5.10**	46.63**	169.69**
Parents	11	73.48**	80.13**	6504.91**	0.91**	26.50**	61.00**	6.32**	15.07**	170.07**
Female	6	50.54**	96.08**	891.96*	0.40**	26.79**	83.30**	7.44**	21.47**	125.04**
Male	4	57.43**	53.43**	217.37	1.89**	32.08**	42.56**	5.38**	8.95	226.64**
Female vs. male	1	275.33**	91.21**	65332.80**	0.03	2.47	0.96	3.39**	1.11	213.99**
Parents vs. hybrids	1	41.46**	124.47**	41908.36**	0.45**	133.51**	283.43**	1.75**	123.42**	39.97*
hybrids	34	71.17**	86.97**	3648.98**	1.05**	39.83**	70.23**	4.81**	54.59**	173.38**
Error	92	1.54	0.96**	333.65	0.02	2.87	2.30	0.16	6.15	7.74

\*, \*\* Significant at P=0.05 and P=0.01 levels respectively.

The mean squares obtained in analysis of variance for combining ability and estimated components of genetic variance for various characters are presented in Table 2.

The result revealed that the mean squares due to females were highly significant for days to flowering, days to maturity, ear head length, ear head girth, test weight and harvest index. This indicated significant contribution of females towards general combining ability variance component for these traits. The hybrids exhibited significant negative general combining ability effects were also a good combiner for early maturity. The results were in accordance with Gill *et al.* [6] and Chavan and Nerkar [5].

The mean squares due to males were significant for days to maturity, ear head length, ear head girth, test weight and grain yield per plant. This indicated the importance of males for their contribution towards

general combining ability variance components. The line × tester mean squares exhibited significance for all plant traits. The ratio of  $\sigma^2_{gca} / \sigma^2_{sca}$  was less than unity for all the characters under study (Table 2). This suggested greater role of non-additive genetic variance in the inheritance of all the traits under study. Among parents, ICMA 07777, ICMA 96222 and 18488 R were good general combiner for grain yield per plant, plant height, number of effective tillers per plant. The female ICMA 96222 was found as good general combiner for characters like grain yield per plant, days to flowering, days to maturity, test weight, harvest index. While ICMA 07777 was good combiner for days to flowering, ear head girth and test weight and harvest index. Among males, 18488-R was found as a good combiner for grain yield per plant, ear head length, ear head girth and test weight (Table 3).

Table 2: Analysis of variance (Mean square) for combining ability, and estimates of components of variance for various characters in pearl millet

Source of variation	df	Days to flowering	Days to maturity	Plant height	No. of effective tillers per plant	Ear head length	Ear head girth	Test weight	Grain yield per plant	Harvest index
Replication	2	4.06	2.6	602.52	0.11 *	2.183	3.64	0.317	6.45	4.16
Crosses	34	71.17 **	86.97 **	3648.98 **	1.04 **	39.82 **	70.23**	4.80**	54.59 **	173.37 **
Females (Line)	6	195.68 **	308.08 **	4683.52	1.41	50.66 *	207.16**	11.10**	71.45	650.57**
Males (Tester)	4	48.62	91.73 *	7414.11	0.42	148.16 **	115.80 *	8.93 *	58.07	44.41
Females × males	24	43.80 **	30.89 **	2762.827 **	1.06 **	19.05 **	28.40 **	2.54**	49.79 **	75.57 **
Error	68	1.43	0.99	440.62	0.02	2.67	2.99	0.18	5.89	9.19
$\sigma^2$ Females		10.12 **	18.47 **	128.05	0.02	2.10 *	11.91 **	0.57 **	1.44	38.33 **
$\sigma^2$ Males		0.22	2.89 *	221.49	-0.03	6.14 **	4.16 *	0.30 *	0.39	-1.48
$\sigma^2_{gca}$		4.35 **	9.38 **	182.55 **	0.01	4.46 **	7.39 **	0.41 **	0.83 *	15.10**
$\sigma^2_{sca}$		14.08 **	9.97 **	809.72 **	0.34 **	5.39**	8.70**	0.79**	14.54 **	22.60 **
$\sigma^2_{gca} / \sigma^2_{sca}$		0.30	0.94	0.22	-0.029	0.82	0.84	0.52	0.05	0.66

\*, \*\* Significant at P=0.05 and P=0.01 levels respectively.

Table 3: Estimation of general combining ability (gca) effects of parents for various characters in pearl millet.

Parents	Days to flowering	Days to maturity	Plant height	No. of effective tillers per plant	Ear head length	Ear head girth	Test weight	Grain yield per plant	Harvest index
ICMA 06777	7.02 **	7.63**	-19.31 **	0.27 **	-2.45 **	3.19 **	-0.09	1.28 *	0.84
ICMA 07777	1.96 **	4.17**	-9.43 *	0.17 **	-0.21	2.59 **	0.38**	1.52 *	-3.17 **
ICMA 96222	-1.77 **	-1.42 **	20.78 **	-0.52 **	3.27**	-0.32	1.36 **	3.46 **	10.99 **
ICMA 97111	-3.63 **	-4.22 **	11.90 *	0.31**	-0.96 *	-2.80 **	0.55 **	-2.81 **	5.30 **
ICMA 98444	-0.77 *	-0.96 **	-6.49	-0.27 **	-0.90 *	0.85 *	-0.32 **	-0.53	-0.84
ICMA 04999	-2.83 **	-5.16 **	-18.61 **	-0.04	1.30 **	-6.81 **	-1.28 **	-1.74**	-9.17**
ICMA 05444	0.02	-0.02	21.17 **	0.06	-0.04	3.29 **	-0.59 **	-1.17	-3.94 **
S.Em. ±	0.64	0.50	9.41	0.08	0.87	0.78	0.21	1.27	1.43
18488 R	0.75 **	2.23**	23.54 **	-0.01	4.35**	0.98**	1.04 **	2.32 **	1.11
18587 R	2.08 **	2.09 **	7.18	0.10 **	0.42	1.54 **	-0.04	0.29	-0.00
18805 R	-1.96 **	-1.47 **	7.28	0.12 **	-0.63	2.47 **	0.06	0.59	0.72
17369 R	-0.67 *	-0.47 *	-22.92 **	-0.23 **	-2.25**	-2.34 **	-0.65 **	-1.81 **	0.67
17548 R	-0.2	-2.38 **	-15.08 **	0.00	-1.89 **	-2.65 **	-0.41 **	-1.39 *	-2.50 **
S.Em. ±	0.54	0.42	7.95	0.07	0.73	0.66	0.17	1.08	1.21

\*, \*\* Significant at P=0.05 and P=0.01 levels respectively.

This suggested that parents might be presumed to have relatively greater number of favorable alleles for developing superior hybrids/varieties of pearl millet (Table 4). Out of 35 hybrids, three hybrids viz., ICMA 07777 × 18488 R (9.53), ICMA 06777 × 18805 R (9.17) and ICMA 96222 × 18587 R (5.42) exhibited

significant positive specific combining ability effects for grain yield per plant. Earlier significant GCA and SCA effects in pear millet for grain yield and component traits also reported by Jagendra Singh *et al.* [8], Patel *et al.* [12], Khandagale *et al.* [10], Priyanka [13] and Mungra *et al* [11].

Table 4: Top three ranking parent with respect to *per se* performance and gca effects and three top ranking hybrids with respect to *per se* performance and sca effects.

Sr. No.	Characters	Best performing parent ( <i>per se</i> performing)	Best combiners	Best performing hybrids ( <i>per se</i> performing)	Status of parents	Hybrids with high sca effects	GCA of the parents	SCA effects
1.	Days to flowering	ICMA 96222	ICMA 97111	ICMA 04999 x 17369 R	G x G	ICMA 07777 x 18805 R	P x G	-5.77
		ICMA 97111	ICMA 04999	ICMA 04999 x 18805 R	G x G	ICMA 05444 x 17548 R	A x A	-5.26
		ICMA 07777	18805 R	ICMA 97111 x 17548 R	G x A	ICMA 98444 x 17548 R	G x A	-5.13
2.	Days to maturity	ICMA 96222	ICMA 04999	ICMA 04999 x 18805 R	G x G	ICMA 07777 x 18805 R	P x G	-6.12
		ICMA 06777	ICMA 97111	ICMA 04999 x 17369 R	G x G	ICMA 97111 x 18587 R	G x P	-3.62
		17548 R	17548 R	ICMA 04999 x 17548 R	G x P	ICMA 05444 x 17369 R	A x G	-3.5
3.	Plant height (cm)	ICMA 07777	17369	ICMA 04999 x 17548 R	G x G	ICMA 06777 x 18587 R	G x A	- 63.66
		ICMA 06777	ICMA 06777	ICMA 06777 x 18587 R	G x A	ICMA 04999 x 17548 R	G x G	- 60.89
		ICMA 98444	ICMA 04999	ICMA 06777 x 17369 R	G x G	ICMA 07777 x 17369 R	G x G	- 38.43
4.	No. of effective tillers/ plant	18805	ICMA 97111	ICMA 97111 x 18587 R	G x P	ICMA 97111 x 18587 R	G x P	1.08
		ICMA 07777	ICMA 06777	ICMA 07777x 18805 R	G x P	ICMA 98444 x 17548 R	P x A	0.94
		18488	ICMA 07777	ICMA 06777 x 17548 R	G x A	ICMA 04999 x 18488 R	A x A	0.70
5.	Ear head length (cm)	18488	18488	ICMA 96222 x 18488 R	G x G	ICMA 06777 x 18805 R	P x A	4.36
		ICMA 07777	ICMA 96222	ICMA 04999 x 18488 R	G x G	ICMA 96222 x 17369 R	G x P	4.10
		17369	ICMA 04999	ICMA 96222 x 18587 R	G x A	ICMA 98444 x 17548 R	P x P	2.80
6.	Ear head girth (mm)	ICMA 07777	ICMA 05444	ICMA 06777 x 18805 R	G x G	ICMA 06777 x 17369 R	G x P	6.07
		18587	ICMA 06777	ICMA 05444 x 18488 R	G x G	ICMA 04999 x 18587 R	P x G	3.42
		18488	ICMA 07777	ICMA 07777 x 18805 R	G x G	ICMA 98444 x 17548 R	G x P	3.29
7.	Test weight (g)	ICMA 07777	ICMA 96222	ICMA 07777 x 18488 R	G x G	ICMA 07777 x 18488 R	G x G	1.50
		17369 R	18488 R	ICMA 96222 x 18587 R	G x A	ICMA 06777 x 18805 R	A x A	1.32
		18488 R	ICMA 97111	ICMA 96222 x 18488 R	G x G	ICMA 96222 x 18587 R	G x A	1.08
8.	Grain yield/ plant (g)	17548 R	ICMA 96222	ICMA 07777 x 18488 R	G x G	ICMA 07777 x 18488 R	G x G	9.53**
		ICMA 96222	18488 R	ICMA 06777 x 18805 R	G x A	ICMA 06777 x 18805 R	G x A	9.17**
		ICMA 98444	ICMA 07777	ICMA 96222 x 18488 R	G x G	ICMA 96222 x 18587 R	G x A	5.42**
9.	Harvest Index (%)	17548 R	ICMA 96222	ICMA 96222 x 17369 R	G x A	ICMA 04999 x 17548 R	P x P	7.17
		ICMA 96222	ICMA 07777	ICMA 96222 x 17548 R	G x P	ICMA 07777 x 17548 R	P x P	6.70
		ICMA 98444	ICMA 06777	ICMA 96222 x 18488 R	G x A	ICMA 98444 x 18805 R	A x A	6.05
		ICMA 05444	ICMA 98444	ICMA 98444 x 17369 R	G x P	ICMA 05444 x 18805 R	G x G	1.54
		ICMA 18587	18805 R	ICMA 97111 x 18587 R	A x A	ICMA 98444 x 17548 R	G x A	1.46

In general all characters of the crosses showing positive and negative combining ability effect, involved either good × good, average × good, good × poor, average × average, average × poor and poor × poor general combiners. Crosses between good × poor and poor × good combiners could be attributed to lack of co-adaptation between favourable alleles of the parents involved. Generally the cross showing high sca effects for grain yield per plant also exhibited high or average or poor sca effects for yield components (Table 4).

This indicated that yield is a complex dependent character which depends on a number of independent component characters.

## CONCLUSIONS

The cross ICMA 07777 × 18488 R showed significant positive specific combining ability effect and high sca effects involving one good general combiner, indicated additive × dominance type of interaction, which can produce desirable transgressive segregate in subsequent generations, also used for exploiting commercial cultivation. The cross ICMA 06777 × 18805 had good × average combiner parents, high *per se* performance and positive specific combining ability effect for grain yield per plant, test weight, ear head length and harvest index. The cross ICMA 96222 × 18488 R had good × good combiner parent, significant positive specific combining effect for grain yield per plant test weight and ear head length. This indicated to obtain desirable transgressive sergeants in F<sub>2</sub> or subsequent generation, may be used for development of pollinated CMS line as female parent and pollen fertility restorer line as male parent for future breeding programme of grain yield per plant, production of high yielding hybrids and for commercial purpose.

## REFERENCES

1. Anand Kumar, K. (1989). Pearl millet: Current status and future potential. *Outlook on Agriculture*, **18**: 46-53.
2. Athwal, D. S. (1965). Hybrid Bajra-1 marks a new era. *Indian Farming*, **15**: 6-7.
3. Brunken, J. N.; DeWet J. M. J. and Harlan J. R. (1977). The morphology and domestication of pearl millet. *Economic Botany*, **31**: 163-174.
4. Burton, G. W. (1958). Cytoplasmic male sterility in pearl millet (*Pennisetum glaucum* (L.) R. Br.). *Agronomy Journal*, **50** (2): 230-231.
5. Chavan, A. A. and Nerkar, Y. S. (1994). Heterosis and combining ability studies for grain yield and its components in pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Journal of Maharashtra Agricultural University*, **19** (1): 58-61.
6. Gill, K. S.; Phul, P. S. and Jindla, L. N. (1974). Line X tester analysis of combining ability in pearl millet [*Pennisetum typhoides* Burn (S. & H.)]. *Journal of Punjab Agricultural University*, **11** (4): 345-349.
7. Hanna, W. W. (1989). Characteristics and stability of a new cytoplasmic-nuclear male-sterile source in pearl millet. *Crop Science*, **29** (4): 1457-1459
8. Jagendra singh and Ravi Sharma (2014). Assessment of Combining Ability in pearl millet Using Line x Tester Analysis. *Advances in Crop Science and Technology*. **2**: 4.
9. Kempthorne, O. (1957). An Introduction to Genetic Statistics. John Willey & Sons. Inc., New York. : 545.
10. Khandagale, S. G.; Vijay Sharma; Lone, R. P.; Khandagale, V. G and Kumara Swamy, R.V. (2014). Combining Ability analysis and gene action in pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Electronic Journal of Plant Breeding*. **5** (3): 445-450.
11. Mungra, K. S.; Dobariya, K. L.; Sapovadiya, M. H and Vavdiya, P. A. (2015). Combining ability and gene action for grain yield and its component traits in pearl millet [*Pennisetum glaucum* (L.) R. Br.]. *Electronic Journal of Plant Breeding*. **6** (1): 66-73.
12. Patel, S. M.; Patel, M. P.; and Patel, J. A. (2014). Combining ability and gene action for grain yield and agronomic traits in pearl millet restorer lines. *Electronic Journal of Plant Breeding*. **5** (3): 395-401.
13. Priyanka, (2015). Heterosis and combining ability analysis for seed yield and its quality traits in pearl millet [*Pennisetum glaucum* (L.) R. Br.]. M.Sc. (Agri.) Thesis (unpublished) submitted to SDAU, Sardarkrushinagar.
14. Rai K. N.; Anand Kumar, K.; Andrews, D. J. and Rao A. S. (2006). Commercial viability of alternative cytoplasmic-nuclear male-sterility systems in pearl millet. *Euphytica*, **121** (3): 107-114.
15. Sukhatme, P.V. and Amble, V.N. (1989). Statistical Methods for Agricultural Workers, ICAR, New Delhi.
16. Suthamathi, P.; Geetha, K. and Sumathi, P. (2007). Combining ability studies on quality characters in pearl millet. *Indian Journal of Genetics*, **67** (2): 203-205.

**Copyright:** © 2019 Society of Education. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.